

***** PROOF OF YOUR ARTICLE ATTACHED, PLEASE READ CAREFULLY *****

After receipt of your corrections your article will be published initially within the online version of the journal.

PLEASE NOTE THAT THE PROMPT RETURN OF YOUR PROOF CORRECTIONS WILL ENSURE THAT THERE ARE NO UNNECESSARY DELAYS IN THE PUBLICATION OF YOUR ARTICLE

READ PROOFS CAREFULLY

ONCE PUBLISHED ONLINE OR IN PRINT IT IS NOT POSSIBLE TO MAKE ANY FURTHER CORRECTIONS TO YOUR ARTICLE

- § This will be your only chance to correct your proof
- § Please note that the volume and page numbers shown on the proofs are for position only

ANSWER ALL QUERIES ON PROOFS (Queries are attached as the last page of your proof.)

- § List all corrections and send back via e-mail to the production contact as detailed in the covering e-mail, or mark all corrections directly on the proofs and send the scanned copy via e-mail. Please do not send corrections by fax or post

CHECK FIGURES AND TABLES CAREFULLY

- § Check sizes, numbering, and orientation of figures
- § All images in the PDF are downsampled (reduced to lower resolution and file size) to facilitate Internet delivery. These images will appear at higher resolution and sharpness in the printed article
- § Review figure legends to ensure that they are complete
- § Check all tables. Review layout, titles, and footnotes

COMPLETE COPYRIGHT TRANSFER AGREEMENT (CTA) if you have not already signed one

- § Please send a scanned signed copy with your proofs by e-mail. **Your article cannot be published unless we have received the signed CTA**

OFFPRINTS

- § 25 complimentary offprints of your article will be dispatched on publication. Please ensure that the correspondence address on your proofs is correct for dispatch of the offprints. If your delivery address has changed, please inform the production contact for the journal – details in the covering e-mail. Please allow six weeks for delivery.

Additional reprint and journal issue purchases

- § Should you wish to purchase a minimum of 100 copies of your article, please visit http://www3.interscience.wiley.com/aboutus/contact_reprint_sales.html
- § To acquire the PDF file of your article or to purchase reprints in smaller quantities, please visit <http://www3.interscience.wiley.com/aboutus/ppv-articleselect.html>. Restrictions apply to the use of reprints and PDF files – if you have a specific query, please contact permreq@wiley.co.uk. Corresponding authors are invited to inform their co-authors of the reprint options available
- § To purchase a copy of the issue in which your article appears, please contact cs-journals@wiley.co.uk upon publication, quoting the article and volume/issue details
- § Please note that regardless of the form in which they are acquired, reprints should not be resold, nor further disseminated in electronic or print form, nor deployed in part or in whole in any marketing, promotional or educational contexts without authorization from Wiley. Permissions requests should be directed to <mailto:permreq@wiley.co.uk>

1

A new numerical method for BLT forward problem based on high-order finite elements

3

Yanbin Hou¹, Jie Tian^{1,2,*},†, Yan Wu¹, Jimin Liang¹ and Xiaowei He¹

5

¹Life Science Research Center, School of Electronics Engineering, Xidian University, Xi'an 710071, Shaanxi, People's Republic of China

7

²Medical Image Processing Group, Institute of Automation, Chinese Academy of Sciences, Beijing 100080, People's Republic of China

9

SUMMARY

Molecular imaging possesses the ability to characterize and measure biological processes at the cellular and molecular level *in vivo*. As one of the new molecular imaging modalities, bioluminescence tomography (BLT) is to reconstruct the light distribution inside a small animal from the photon flux measured on its surface. To obtain accurate and robust reconstruction, it needs a good understanding of the propagation of photons in biological tissues, which is referred to as the forward problem in BLT. Because the turbid media is high scattering and low absorption, the propagation process can be described by the steady diffusion equation. In this paper, an *hp-adaptivity* method for BLT forward problem is presented based on finite elements of high orders and moderate meshes by finite element method (FEM). Both numerical simulation and physical experiment are performed to evaluate the accuracy of solution and the efficiency of computation. The relevant results show that element order is more critical than mesh size to produce an accurate FEM solution efficiently. Copyright © 2008 John Wiley & Sons, Ltd.

Received 1 April 2008; Revised 20 October 2008; Accepted 22 October 2008

KEY WORDS: molecular imaging; bioluminescence tomography; forward problem; high-order finite elements; diffusion approximation

23

1. INTRODUCTION

Modern medical imaging has undergone structural and functional imaging techniques since the discovery of X-ray by Wilhelm Konrad Roentgen in 1895. The relevant equipments, such as CT,

25

*Correspondence to: Jie Tian, Medical Image Processing Group, Institute of Automation, Chinese Academy of Sciences, Beijing 100080, People's Republic of China.

†E-mail: tian@IEEE.org

Contract/grant sponsor: National Basic Research Program of China (NBRPC); contract/grant number: 2006CB705700

Contract/grant sponsor: Program for Cheung Kong Scholars and Innovative Research Team in University (PCSIRT); contract/grant number: IRT0645

Contract/grant sponsor: Chair Professors of Cheung Kong Scholars Programme

Contract/grant sponsor: Shaanxi Science and Technology Programming; contract/grant number: 2007K08-06

1 MRI, PET, SPECT, etc, have brought great innovation in the diagnosis and treatment of diseases.
2 These medical equipments help medical staff to observe and measure the tissue lesions more
3 clearly and accurately. However, most of them are only able to screen the advanced stages of
4 diseases. With the advances in biomedical and genetic engineering, molecular imaging provides
5 promising solutions to early diagnosis of disease, personalized treatment, drug development, gene
6 study and so on [1–3]. Compared with other molecular imaging techniques, optical imaging
7 has the advantages of nano–molar sensitivity, high spatial resolution, short imaging time, low
8 cost and so on [4, 5]. Optical imaging modalities include optical coherent tomography, diffusion
9 optical tomography, photoacoustic tomography, optical-PET, bioluminescence tomography (BLT)
10 and fluorescence molecular tomography [6, 7]. BLT is to reconstruct the bioluminescent light
11 sources inside an object from the optical signals measured at its boundary, and to provide real-time
12 detection and quantitative measurement of tagged cells while keeping the small living animals
13 intact. Although optical imaging can only detect the surface light, BLT fails to get the depth
14 information of targets under test directly [8]. Multi-spectral and multi-modality method are recently
15 developed to improve the performance of BLT [9–11].

16 Since biological tissues are turbid, bioluminescent photons undergo absorption, reflection, scatter-
17 ing and other optical behaviors. BLT employs the near infrared light to image *in vivo* because
18 the lowest coefficient of absorption occurs within the range of 650–900 nm [3]. Light propagation
19 in biological tissues can be depicted by Monte Carlo method (MC) based on repeated random
20 sampling, or by radiative transfer equation (RTE) based on classic transport theory [6, 12–14].
21 But both methods are computationally expensive. In high-scattering and low-absorption biological
22 tissues, RTE is simplified as the diffusion equation (DE) with appropriate boundary condition,
23 which provides a quite accurate description of the imaging model [15]. BLT problems consist of
24 forward problem and inverse problem. Forward problem is to find out the light distribution with
25 given light sources and tissue parameters, whereas inverse problem is either to determine tissue
26 optical properties or to retrieve inner light sources from boundary measurement [16]. Mathemati-
27 cally, forward problem is to solve a partial differential equation. The solution is difficult to obtain
28 analytically in most practical cases. Many numerical methods, such as finite difference methods,
29 finite element method (FEM), boundary element method, are widely adopted [17–21]. Owing to
30 the efficient and flexible discretization techniques, FEM has become well established in biomedical
31 optics [22].

32 In this paper, we investigate the effects of mesh size and element order on FEM solution and
33 evaluate the accuracy and the convergence compared with analytical solution, MC simulation and
34 experimental measurement. In the second section, BLT forward problem is formulated by DE
35 and then analyzed via FEM. In the third section, numerical simulation and physical experiment
36 are performed to evaluate the accuracy and the efficiency of FEM. In the final section, results and
37 conclusions are discussed.

38 2. THEORY AND METHOD

39 2.1. Transport theory and RTE

In BLT, forward problem is to find the photon density in the biological tissue and the outgoing
flux on its boundary. Photon transport in biological tissue can be described with RTE based on

transport theory [12, 19]

$$\frac{\partial L(r, \hat{s}, t)/V}{\partial t} = -\hat{s} \cdot \nabla L(r, \hat{s}, t) - (\mu_a + \mu_s)L(r, \hat{s}, t) + \mu_s \int_{4\pi} L(r, \hat{s}', t) P(\hat{s}' \cdot \hat{s}) d\hat{s}' + S(r, \hat{s}, t) \quad \forall r \in \Omega \quad (1)$$

- 1 where $L(r, \hat{s}, t)$ denotes radiance at position r at time t into direction \hat{s} in the unit of $\text{Wmm}^{-2} \text{sr}^{-1}$,
 3 V the velocity of light in tissues, μ_a and μ_s are absorption and scattering coefficients, respectively,
 in mm^{-1} , Phase function $P(\hat{s}' \cdot \hat{s})$ represents the probability distribution of light with propagation
 direction \hat{s}' scattered into the direction \hat{s} , $S(r, \hat{s}, t)$ is the inner light source in $\text{Wmm}^{-3} \text{sr}^{-1}$.

5 **2.2. Diffusion approximation and boundary conditions**

- 7 The RTE is difficult to solve since there are six independent variables. The RTE is usually simplified
 by diffusion approximation for high-scattering and low-absorption medium ($\mu_a \ll \mu_s$), where the
 radiance $L(r, \hat{s}, t)$ is expanded to the first order of spherical harmonics. Thus the RTE (1) can be
 9 reduced to the DE [16]

$$\nabla \cdot D(\nabla \Phi(r, t)) - \mu_a \Phi(r, t) = \frac{\partial \Phi(r, t)/V}{\partial t} - S(r, t) \quad \forall r \in \Omega \quad (2)$$

- 11 where $\Phi(r, t)$ is the instantaneous photon flux density at time t in Wmm^{-2} , $D = 1/(3(\mu_a + (1 -$
 $g)\mu_s))$ the optical diffusion coefficient in mm and g the anisotropy parameter.
 13 Let $S(r, t) = \delta(r, t)$ and Ω be an infinite homogeneous domain, we can get Green's function
 solution to the DE:

$$15 \quad \Gamma(r, t) = V(4\pi DVt)^{-3/2} \exp[-r^2/(4DVt) - \mu_a Vt] \quad \forall r \in \Omega \quad (3)$$

For an arbitrary source, the solution to Equation (2) can be obtained by the following convolution:

$$17 \quad \Phi(r, t) = \Gamma(r, t) * S(r, t) = \int_0^t \int_{\Omega} \Gamma(r - r', t - t') S(r', t') dr' dt' \quad (4)$$

- 19 If the measurement is carried out in an ideal environment without outer light interference, the
 boundary condition for Equation (2) can be expressed as [23]

$$\Phi(r, t) + 2A(r; n, n') D(v \cdot \nabla \Phi(r, t)) = 0 \quad \forall r \in \partial\Omega \quad (5)$$

- 21 where v is the unit outer normal on boundary, n and n' are the refractive indices of inner and outer
 medium, respectively, $A(r; n, n')$ can be approximated as

$$23 \quad A(r; n, n') \approx (1 + R(r))/(1 - R(r)) \quad (6)$$

- where n' is close to 1 when experiments are performed in air, $R(r)$ is a parameter governing the
 25 internal reflection on boundary $\partial\Omega$ [19]:

$$R(r) \approx -1.4399n^{-2} + 0.7099n^{-1} + 0.6681 + 0.0636n$$

- 27 According to Fick's law, the outgoing flux density measured on the boundary is [8]:

$$G(r, t) = -D(v \cdot \nabla \Phi(r, t)) = \Phi(r, t)/(2A(r; n, n')) \quad (7)$$

1 In real experiments, the bioluminescent sources are usually provided by mixing different chem-
 2 ical solutions, which can release constant power during quite a long period. This means we usually
 3 deal with photon flux density independent of time. From Equations (2) and (5), we can get the
 steady-state form of DE and its boundary condition as follows:

$$\begin{aligned}
 & -\nabla \cdot (D \nabla \Phi(r)) + \mu_a \Phi(r) = S(r) \quad \forall r \in \Omega \\
 & \Phi(r) + 2A(r; n, n') D(v \cdot \nabla \Phi(r)) = 0 \quad \forall r \in \partial\Omega
 \end{aligned} \tag{8}$$

2.3. Analytical solution to steady-state DE

7 It is difficult to obtain an analytical solution to the DE in most practical cases. However, when
 the model is simple, for example, infinite homogeneous medium enclosing a spherical source,
 9 the analytical solution can be calculated through Equation (4). The power density of a uniform
 spherical solid source in a three-dimensional spherical coordinate system can be written as [16, 24]

$$S(r, \theta, \phi) = \begin{cases} p & \text{for } 0 \leq \theta \leq 2\pi, 0 \leq \phi \leq \pi, 0 \leq r \leq r_0 \\ 0 & \text{else} \end{cases} \tag{9}$$

11 where p is the light source power in W , r_0 is the radius of the spherical source in mm, θ denotes
 13 the polar angle and ϕ the azimuthal angle.

15 Substituting Equation (9) into Equation (4) and simplifying the convolution, we can derive the
 analytical solution to the steady-state DE [24]:

$$\Phi(r) = \frac{p \exp[-\mu_{\text{eff}} r]}{r D (\mu_{\text{eff}})^2} \left(r_0 \cosh(\mu_{\text{eff}} r_0) - \frac{1}{\mu_{\text{eff}}} \sinh(\mu_{\text{eff}} r_0) \right) \tag{10}$$

17 where r_0 denotes the radius of the spherical source, r denotes the distance between the source
 surface and measuring point, μ_{eff} is the effective attenuation coefficient defined as $\mu_{\text{eff}} = \sqrt{\mu_a / D}$.

19 2.4. Weak form and FEM formulation

The weak form of Equation (8) is to find $\Phi(r)$ such that

$$\int_{\Omega} D \nabla \Phi(r) \cdot \nabla \Psi(r) \, dr + \int_{\Omega} \mu_a \Phi(r) \Psi(r) \, dr + \int_{\partial\Omega} \frac{1}{2A} \Phi(r) \Psi(r) \, dr = \int_{\Omega} S(r) \Psi(r) \, dr \tag{11}$$

21 where $\Psi(r)$ is an arbitrary piecewise continuous test function [23]. According to the standard finite
 23 element analysis, domain Ω is decomposed into T vertex nodes (N_1, N_2, \dots, N_T) and N_e mesh
 elements, denoted as e_l ($l = 1, 2, \dots, N_e$). Let the set $E = \{e_l\}$ be triangulation. Then $\Phi(r)$ can be
 25 expressed approximately as

$$\Phi(r) \approx \Phi^h(r) = \sum_{k=1}^T \phi_k \varphi_k(r) \quad \forall r \in \Omega \tag{12}$$

27 where ϕ_k is the approximate nodal value of $\Phi(r)$ on the node N_k , $\varphi_k(r)$ is the nodal basis function
 with support over the elements e_l , which have the node N_k as a common vertex. In the same way,
 29 the source is approximated as

$$S(r) \approx S^h(r) = \sum_{k=1}^{T_s} s_k \gamma_k(r) \quad \forall r \in \Omega \tag{13}$$

1 where $\{s_k\}$ is the interpolation node values of $S(r)$ and $\gamma_k(r)$ the interpolation basis functions,
 3 which may be the same with or different from $\varphi_k(r)$, T_s is the number of interpolation basis
 functions. Usually, tissue consists of several organs and T_s is much smaller than T . We adopt a
 5 global node numbering scheme for each organ and light source to deal with the discontinuities in
 optical coefficients. Using $\varphi_k(r)$ as the test function and substituting Equations (12) and (13) into
 Equation (11), we have:

$$7 \quad \sum_{i,j} \left\{ \int_{\Omega} D \nabla \varphi_i \cdot \nabla \varphi_j \, dr + \int_{\Omega} \mu_a \varphi_i \varphi_j \, dr + \int_{\partial\Omega} \frac{1}{2A} \varphi_i \varphi_j \, dr \right\} \phi_i = \sum_{i,j} \left\{ \int_{\Omega} \gamma_i \varphi_j \, dr \right\} s_i \quad (14)$$

By defining the system matrix $M = (M_{ij}) \in R^{T \times T}$ and the vector $F = (f_i) \in R^{T \times 1}$:

$$M_{ij} = \int_{\Omega} D \nabla \varphi_i \cdot \nabla \varphi_j \, dr + \int_{\Omega} \mu_a \varphi_i \varphi_j \, dr + \int_{\partial\Omega} \frac{1}{2A} \varphi_i \varphi_j \, dr$$

$$f_i = \sum_j \left\{ \int_{\Omega} \gamma_i \varphi_j \, dr \right\} s_j$$

Equation (14) becomes a linear equation as follows:

$$9 \quad M \phi = F \quad (15)$$

where $\phi = (\phi_i)$, $i = 1, 2, \dots, N_e$. M is a sparse, symmetric and positive-definite system matrix.
 11 Hence, the solution to Equation (8) is to find out a series of nodal values of $\Phi(r)$.

As a linear equation, Equation (15) is generally solved by direct solvers or iterative solvers. The
 13 formers introduce matrix factorization and Gauss elimination, while the latters need initial values
 and proper preconditioners to accelerate the convergence.

15 2.5. High-order basis functions

As was mentioned above, domain under investigation is partitioned into a set of mesh elements.
 17 These finite elements are of simple shapes, such as triangle, quadrilateral for 2D domain and
 tetrahedron, prism for 3D domain. Since triangular elements have better approximation of irreg-
 19 ular geometry, we employ tetrahedral elements for subdomains and triangular elements for the
 boundary.

21 After the discretization of domain Ω into finite elements, approximations to the dependent
 variables need to be introduced, as illustrated in Equation (12). That means one should approximate
 23 $\Phi(r)$ with a function $\Phi^h(r)$ in finite element spaces. ϕ_k is called *degrees of freedom* (DOFs),
 and $\varphi_k(r)$ the nodal basis function. In terms of the finite element, basis function can be linear,
 25 quadratic, cubic or even higher order [25, 26]. In this paper, we construct piecewise continuous
 basis functions by using Lagrange interpolation.

27 For 1D domain, $\Phi(x)$, which passes through $n + 1$ points $(x_0, y_0), (x_1, y_1), \dots, (x_n, y_n)$, is a
 linear combination of Lagrange interpolation basis functions:

$$\Phi(x) = \sum_{k=0}^n y_k N_k(x)$$

$$N_k(x) := \prod_{\substack{j=0 \\ j \neq k}}^n \frac{(x - x_j)}{(x_k - x_j)}$$

29

1 It is obvious that 1D Lagrange basis functions satisfy following properties [27]:

- 2 (1) $0 \leq N_k(x) \leq 1$.
- 3 (2) $N_k(x_j) = \delta_{jk}$.
- 4 (3) $\sum_k N_k(x) = 1$.

5 According to the definition in Equation (16), $\varphi_k(x)$ is of degree at most $n - 1$. The more points that are used in the interpolation, the higher the degree of the basis function, as shown in Figure 1.

7 For 2D domain, the description of basis functions is simplified by the introduction of *barycentric coordinates* [28] for simplicity. In the context of triangle, barycentric coordinates are also known as area coordinates. As illustrated in Figure 2, coordinates of P with respect to triangle ABC are proportional to the areas of PBC , PCA and PAB .

11 If we define:

$$L_1 = \frac{S_{\Delta PBC}}{S_{\Delta ABC}}, \quad L_2 = \frac{S_{\Delta PCA}}{S_{\Delta ABC}}, \quad L_3 = \frac{S_{\Delta PAB}}{S_{\Delta ABC}} \quad (17)$$

13 The area coordinates (L_1, L_2, L_3) of vertices A, B and C are $(1, 0, 0)$, $(0, 1, 0)$ and $(0, 0, 1)$, respectively. Obviously, L_1, L_2 and L_3 satisfy properties (1), (2) and (3). Thus we can express




Figure 1. 1D Lagrange interpolation basis function: (a) linear; (b) quadratic; and (c) cubic.




Figure 2. Triangular area coordinates for linear basis function.




Figure 3. Triangular area coordinates for quadratic basis function.

1 $\Phi(r)$ as a combination of linear Lagrange interpolation basis functions

$$\Phi(r) = \sum_{k=1}^3 \phi_k \varphi_k(r) \quad \forall r \in \Delta P_i P_j P_m \quad (18)$$

3 where $\varphi_k = L_k, k = 1, 2, 3$.

5 Because area coordinates do not depend neither on the shape nor on the position of the triangular finite element, it is quite convenient to derive the quadratic and cubic Lagrange interpolation basis functions

$$\varphi_k = \prod_{j=1}^m \frac{f_j^{(k)}(L_1, L_2, L_3)}{f_j^{(k)}(L_{1k}, L_{2k}, L_{3k})}, \quad k = 1, 2, \dots, N \quad (19)$$

7 where m is the degree of $\varphi_k, f_j^{(k)}(L_1, L_2, L_3)$ is the left-hand side of straight-line equation excluding the k th node, $f_j^{(k)}(L_{1k}, L_{2k}, L_{3k})$ the area coordinates on the k th node, $N = (m+1)(m+2)/2$ the number of nodes contained in the triangular element.

11 Based on Equation (19), it needs to add other three nodes to build quadratic basis function, as shown in Figure 3. Nodes 4 to 6 are the midpoints of three edges, and $\Phi(r)$ is expressed as follows:

$$\Phi(r) = \sum_{k=1}^6 \phi_k \varphi_k(r) \quad \forall r \in \Delta ABC \quad (20)$$

15 where $\varphi_k = L_k(2L_k - 1), k = 1, 2, 3, \varphi_4 = 4L_2L_3, \varphi_5 = 4L_3L_1, \varphi_6 = 4L_1L_2$.

17 Likewise, the basis functions of different orders for 3D can be constructed by the introduction of volume coordinates. In general, basis function of high order specifies a finite element space with better accuracy while the drawback is that DOFs increase.

19 2.6. Error estimation and convergence

21 FEM formulation is the discretization of the original problem, the error between exact solution and approximation is bounded by [29]

$$\|\Phi - \Phi^h\| \leq Ch^{\min(m-1, p)} \quad (21)$$

1 where $\|\cdot\|$ denotes the Euclidean norm, h is the maximum mesh size, p is the degree of the basis
 3 function and m is a measure of the order of the singularity in the problem. In order to reduce the
 5 error, three versions of FEM may be adopted, which correspond to shortening the element size by
h-adaptivity, increasing the degree of basis function by *p-adaptivity* and swapping the edge and
 facet of the element by *r-adaptivity*, respectively.

7 In theory, if the maximum mesh size is sufficiently small, the numerical solution to Equation (15)
 9 could converge to the exact solution of Equation (8) [23]. But the smaller h is, the more unknowns
 11 in Equation (15), need to be found out. In practice, we usually have to make a tradeoff between the
 solution accuracy and computation overhead. In this paper, *h-p* FEM is applied, where convergence
 can be achieved by refining mesh element or by increasing the approximation order or by a
 combination of both [30]. Convergence order n is evaluated by:

$$n = \frac{\ln \|\Phi - \Phi^h\| - \ln(c)}{\ln(h)}, \quad c \in (0, 1] \quad (22)$$

13 In the next section, we will study the effects of h and p on the accuracy and the convergence
 of FEM solutions with numerical simulation and physical experiment.

15 3. SIMULATION AND EXPERIMENT

In study of BLT forward problem by FEM, the domain under consideration is first described
 17 by creating geometry. Second, the region of interest (ROI) is discretized into finite elements by
 19 meshing, and then through assigning relevant parameters and selecting appropriate basis functions
 we form a group of linear equations as described in Section 2. Finally, a proper solver is employed
 21 and the unknown variables are determined. Extensive comparisons were made to evaluate the
 effects of high-order FEM on the accuracy of solution and the efficiency of computation. In our
 23 simulations, we partitioned the geometric model into tetrahedral elements with Netgen 4.3, and the
 main computational platform is a Compaq desktop computer, dc7700 with Intel Core™ 2 CPU
 6300 @ 1.86 GHz and 2 GB RAM.

25 3.1. Comparison with analytical solution

3.1.1. *The accuracy of FEM solution.* In order to examine the accuracy compared with the analyt-
 27 ical solution obtained by Equation (10), we adopted the homogeneous phantom from Hui [16].
 As shown in Figure 4(a), the phantom consists of concentric spheres. The inner ball of the radius
 29 0.3 mm was a solid uniform light source with power of 1 μ W. The outer was the tissue with optical
 parameters of $\mu_a = 0.082 \text{ mm}^{-1}$, $\mu_s = 10.27 \text{ mm}^{-1}$, $g = 0.90$ and $n = 1.37$. Figure 4 (b) shows the
 31 phantom mesh after discretization with a radius of 4 mm and moderate mesh size. With radius
 of tissues varying from 2 to 10 mm, we obtained a group of flux density values. Because the
 33 solutions to Equation (15) were the nodal values of $\Phi(r)$ on overall nodes, whereas the solution to
 Equation (10) was a point value on the phantom surface, some work must be done as a necessary
 35 preprocessing step. Here, we integrated solutions over the tissue surface to derive the photon flux
 in Watt.

37 As it was expected, both results from FEM and analytical formula declines with the increasing
 radius of phantom surface, as shown in Figure 5. The numerical solutions show good accordance
 39 with the analytical solutions.




Figure 4. Spherical homogenous phantom containing a solid ball source: (a) geometric model and (b) mesh elements after moderate discretization.




Figure 5. Comparison between FEM solution and analytical solution.

1 *3.1.2. The convergence of high-order elements.* We constructed basis functions by Lagrange inter-
 2 polation in this paper. The basis function is a piecewise continuous on each element and its degree
 3 is equal to the order of the Lagrange element. The higher the order is, the smoother the basis
 4 function is, and the better the numerical solution approaches the true solution [15, 24]. Three
 5 groups of FEM solutions were obtained by adopting tetrahedral elements of different orders, all
 6 the geometric and optical parameters were the same as those in Section 3.1.1 except that the tissue
 7 was fixed with a radius of 4 mm. Figure 6 demonstrates the comparison, and Tables I and II list
 8 the quantitative results.

9 As shown in Figure 6, fast convergence can be achieved, when the exact solution is smooth and
 10 higher-order finite elements are adopted. Particularly, the FEM solution with cubic element can
 11 achieve a fairly good consistency with the analytical solution even on coarsest meshes. Hence, it
 12 is quite wise to employ elements of higher order than just to refine the meshes further, especially
 13 when both of them promise the similar accuracy, as listed in Tables I and II. For BLT forward
 14 problem, FEM solution with quadratic element on coarse mesh is able to make a good tradeoff
 15 between precision and efficiency.

3.2. Comparison with MC simulation

17 Because of its accuracy and flexibility, MC method has been established as a gold standard for
 photon propagation simulation in turbid medium. We also performed the numerical experiment to




Figure 6. FEM solutions with tetrahedral elements of different orders.

Table I. The quantitative results of FEM solutions with different order elements compared with the analytical solution, which are considered as exact solution and its power is 0.236 mW.

Finite element	Tetrahedra	Degree of freedoms	Relative error	Elapsed time
Linear	226 599	40 214	-0.85%	13.5 s
Quadratic	2124	2058	-0.85%	0.6 s
Cubic	1158	5062	+0.85%	1.2 s

Table II. The convergence orders of FEM solutions with different order elements.

Mesh sizes	Tetrahedra	Linear element		Quadratic element		Cubic element	
		Error norm	Convergence order	Error norm	Convergence order	Error norm	Convergence order
Coarsest	1158	0.062	0.5	0.003	3.6	0.002	4.1
Coarser	2124	0.039	0.8	0.002	3.4	0.002	3.4
Coarse	4456	0.014	1.4	0.001	3.3	0.002	2.8
Moderate	11070	0.013	1.2	0.001	2.7	0.002	2.3
Fine	20847	0.009	1.2	0.001	2.4	0.002	2.0
Finer	58798	0.005	1.3	0.002	1.7	0.002	1.7
Finest	220001	0.002	1.4	0.002	1.4	0.002	1.4

1 compare the accuracy between MC and quadratic FEM. A cylindrical phantom of diameter 19 mm
 and height 30 mm was used. It had the optical parameters of $\mu_a = 0.01 \text{ mm}^{-1}$, $\mu_s = 3.0 \text{ mm}^{-1}$,
 3 $g = 0.80$ and $n = 1.37$. The phantom was discretized into 20 365 tetrahedral elements and 3805
 nodes. A total 301 virtual detectors were placed on the phantom surface to record the photon
 5 flux density. The phantom was embedded with two light sources, whose parameters were listed in
 Table III.

Table III. The parameters of light source.

Source no.	Shape	Center (mm)	Diameter (mm)	Height (mm)	Power (nW)
1	Cylinder	(0, 0, 19.9)	0.4	1	170
2	Cylinder	(6, 0, 19.9)	0.4	1	170




Figure 7. Comparison between FEM solution and MC simulation.

1 It took about 2 h to perform MC simulation in molecular optical simulation environment [31].
 2 Figure 7 shows that the solution obtained by quadratic FEM is in good agreement with the result
 3 from MC simulation with a relative error (RE) of 6.3%. The RE is defined as $\sum_{\text{detectors}} |\Phi_{\text{MC}} - \Phi_{\text{FEM}}| / \sum_{\text{detectors}} \Phi_{\text{MC}}$ [32]. Compared with MC, the quadratic FEM had an excellent computation
 5 efficiency that it spent less than 8 s in solving the same problem.

3.3. Comparison with CCD measurement

7 We adopted the same phantom and method to perform the physical experiment, as reported in the
 8 literatures [23, 33]. The mouse chest phantom of 30 mm height and 30 mm diameter consisting of
 9 four different high molecular polymers to represent muscle (M), lungs (L), heart (H) and bone
 (B), respectively, as shown in Figure 8(a). Table IV lists the corresponding optical parameters.

11 The luminescent light stick (Glowproducts, Canada) was selected as the testing light source,
 which could last for about 4 h at an emission wavelength around 700 nm. Two light tubes were
 13 placed inside the two small holes with their centers at $(-9.0, 1.5, 15.0)$ and $(-9.0, -1.5, 15.0)$
 in the left lung. Their power were 105.1 and 97.4 nW, respectively. A scientific nitrogen-cooled
 15 CCD camera (Roper Scientific Inc, Trenton, NJ) was used for recording the flux density on the
 cylindrical surface of the phantom, as schematically shown in Figure 8(b).

17 While obtaining FEM solutions, we had implemented three groups of simulation under different
 settings. The phantom was discretized into 21 673 linear elements with the coarsest mesh size,
 19 65 529 linear elements with coarse mesh size and 21 422 quadratic elements with the coarsest mesh
 size, in Simulation 1, 2 and 3, respectively. Figure 9 gives the CCD measurement and the result
 21 of Simulation 3. The image in Figure 9(b) was transformed from the flux density on the phantom




Figure 8. Heterogeneous physical phantom: (a) the phantom consisting of muscle (M), lungs (L), heart (H) and bone (B) and (b) a cross-section through light sources in the left lung. The yellow arrow shows the camera direction during data acquisition.

Table IV. Optical parameters of the physical phantom

Material	μ_a (mm^{-1})	μ_s (mm^{-1})	g	n
Tissue (T)	0.007	10.31	0.90	1.37
Lung (L)	0.023	20.00	0.90	1.00
Heart (H)	0.011	1.96	0.90	1.37
Bone (B)	0.010	3.00	0.98	1.37




Figure 9. Bioluminescent images of front view: (a) CCD measurement and (b) FEM simulation.

1 surface by geometric mapping. Both CCD measurement and FEM simulation data were mapped
 into 24 b bitmap after regularization to simplify the comparison.

3 We sampled the middle two columns in Figures 9(a) and (b) to make quantitative comparisons,
 and detailed results are plotted in Figure 10.

5 As illustrated in Figure 10, simulation 3 with quadratic elements and the coarsest meshes
 matches CCD measurement best. The mean REs of simulation 1, 2 and 3 are 4.2, 3.2 and 2.8% and
 7 the maximum REs are 20.3, 14.3 and 6.8%, respectively. It took 6.9, 15.7 and 12.6 s to perform




Figure 10. Relative errors (RE) between experimental measurement and three groups of numerical simulation.

1 simulation 1, 2 and 3. That means simulation 3 based on quadratic elements has the obvious
 2 advantages of accuracy and efficiency over simulation 1 and 2, which are based on linear elements.

3

4. DISCUSSIONS AND CONCLUSIONS

5 FEM is feasible for the numerical study of BLT forward problem. In this paper, we elucidate the
 6 BLT forward problem, introduce the formulation by finite element analysis and discuss FEMs with
 7 varying mesh size and varying basis function degree. Both numerical and experimental studies
 8 on the accuracy and the convergence of FEM solution are made. As the results indicate, based
 9 on proper mesh size and basis function degree, accurate FEM solutions to DE can be obtained
 10 efficiently. Moreover, adopting higher-order finite elements will achieve much better accuracy
 11 and convergence than just refining the meshes further. For BLT forward problem, quadratic finite
 12 elements of coarse size are able to make a good tradeoff between solution accuracy and computation
 13 efficiency.

14 Since the system matrix is sparse and symmetric, specialized techniques on the storage and
 15 manipulation for sparse matrix can be employed. When decomposing the geometry into mesh
 16 elements, we can only refine ROI (e.g. light source domain), while keeping the remainder intact.
 17 Our future work is to focus on the adaptive meshing and multiple sources with complicated tissues.
 18 Relevant results will be available later.

19

ACKNOWLEDGEMENTS

This work is supported by the Chair Professors of Cheung Kong Scholars Programme, the Program
 for Cheung Kong Scholars, Innovative Research Team in University (PCSIRT, IRT0645), National Basic

Research Program of China (NBRPC, 2006CB705700), and Shaanxi Science and Technology Programming (2007K08-06).

1

REFERENCES

1. Weissleder R, Mahmood U. Molecular imaging. *Radiology* 2001; **219**:316–333.
- 3 2. Dzik-Jurasz ASK. Molecular imaging in vivo: an introduction. *British Journal of Radiology* 2003; **76**:S98–S109.
- 5 3. Cherry SR. In vivo molecular and genomic imaging: new challenges for imaging physics. *Physics in Medicine and Biology* 2004; **49**:R13–R48.
- 7 4. Massoud TF, Gambhir SS. MI in living subjects: seeing fundamental biological process in a new light. *Genes and Development* 2003; **17**(5):545–580.
- 9 5. Ntziachristos V, Ripoll J, Wang LV, Weissleder R. Looking and listening to light: the evolution of whole-body photonic imaging. *Nature Biotechnology* 2005; **23**(3):313–320.
- 11 6. Wang LV, Wu H. *Biomedical Optics: Principles and Imaging*. Wiley: New York, 2007.
- 13 7. Wang G, Jiang M, Tian J, Cong W, Li Y, Han W, Kumar D, Qian X, Shen H, Zhou T, Cheng J, Lv Y, Li H, Luo J. Recent development in bioluminescence tomography. *Current Medical Imaging Reviews* 2006; **2**:453–457.
- 15 8. Yujie LV, Tian J, Cong W, Wang Ge, Luo J, Yang W, Li H. A multilevel adaptive finite element algorithm for bioluminescence tomography. *Optics Express* 2006; **14**(18):8211–8223.
- 17 9. Cong A, Wang G. Multispectral bioluminescence tomography: methodology and simulation. *International Journal of Biomedical Imaging* 2006; **2006**:1–7. DOI:10.1155/IJBI/2006/57614.
- 19 10. Wang G, Shen H, Kumar D, Qian X, Cong W. The first bioluminescence tomography system for simultaneous acquisition of multi-view and multi-spectral data. *International Journal of Biomedical Imaging* 2006; **2006**:1–8. DOI:10.1155/IJBI/2006/58601.
- 21 11. Lv Y, Tian J, Cong W, Wang Ge, Yang W, Qin C, Xu M. Spectrally resolved bioluminescence tomography with adaptive finite element analysis: methodology and simulation. *Physics in Medicine Biology* 2007; **52**(15):4497–4512.
- 23 12. Ishimaru A. *Wave Propagation and Scattering in Random Media*. Academic Press: New York, 1978.
- 25 13. Li H, Tian J, Zhu F, Cong W, Wang LV, Hoffman EA, Wang Ge. A mouse optical simulation environment (MOSE) to investigate bioluminescent phenomena in the living mouse with the Monte Carlo method. *Academic Radiology* 2004; **11**(9):1029–1038.
- 27 14. Wang LH, Jacques SL, Zheng LQ. MCML-Monte Carlo modeling of photon transport in multi-layered tissues. *Computer Methods and Programs in Biomedicine* 1995; **47**:131–146.
- 29 15. Cong WX, Kumar D, Liu Y, Cong A, Wang G. A practical method to determine the light source distribution in bioluminescent imaging. *Proceedings of the SPIE* 2004; **5535**:679–686.
- 31 16. Hui L. Studies on forward and inverse problems in in vivo bioluminescent tomography imaging. *Ph.D. Dissertation*, Institute of Automation, Chinese Academy of Sciences, Beijing, 2005.
- 33 17. Liu J-L. A finite difference method for symmetric positive differential equation. *Mathematical Computations* 1994; **62**:105–118.
- 35 18. Arridge SR, Schweiger M, Hiraoka M, Delpy DT. A finite element approach for modeling photon transport in tissue. *Medical Physics* 1993; **20**:299–309.
- 37 19. Schweiger M, Arridge SR, Hiraoka M *et al.* The finite element method for the propagation of light in scattering media: boundary and source condition. *Medical Physics* 1995; **22**:1779–1792.
- 39 20. Cong W, Wang G. Boundary integral method for bioluminescence tomography. *Journal of Biomedical Optics* 2006; **11**(2):020503.
- 41 21. Song CM, Bazyar MH. Development of a fundamental-solution-less boundary element method for exterior wave problems. *Communications in Numerical Methods in Engineering* 2008; **24**(4):257–279.
- 43 22. Arridge SR, Dehghani H, Schweiger M *et al.* The finite element model for the propagation of light in scattering media: a direct method for domains with nonscattering regions. *Medical Physics* 2000; **27**:252–264.
- 45 23. Cong W, Wang G, Kumar D, Liu Y, Jiang M, Wang LV, Hoffman EA, McLennan G, McCray PB, Zabner J, Cong A. Practical reconstruction method for bioluminescence tomography. *Optics Express* 2005; **13**:6756–6771.
- 47 24. Cong W, Wang LH, Wang G. Formulation of photon diffusion from spherical bioluminescent sources in an infinite homogeneous medium. *Biomed Engineering Online* 2004.
- 49 25. Kuzmin D. A high-resolution finite element scheme for convection-dominated transport. *Communications in Numerical Methods in Engineering* 2000; **16**(3):215–223.

Q1

- 1 26. Lu H, Berzins M, Goodyer CE, Jimack PK. High-order discontinuous Galerkin method for elastohydrodynamic
lubrication line contact problems. *Communications in Numerical Methods in Engineering* 2005; **21**(11):643–650.
- 3 27. Idelsohn SR, Onate E, Calvo N, Pin FD. The meshless finite element method. *International Journal for Numerical
Methods in Engineering* 2003; **58**(6):893–912.
- 5 28. Schöberl J, Zaglmayr S. High order Nédélec elements with local complete sequence properties. *COMPEL: The
International Journal for Computation and Mathematics in Electrical and Electric Engineering* 2005; **24**(2):
7 374–384.
- 9 29. Yujie Lv, Tian J, Li H, Luo J, Cong W, Wang Ge, Kumar D. Modeling the forward problem based on the adaptive
FEMs framework in bioluminescence tomography. *Proceedings of SPIE, Developments in X-Ray Tomography V*,
2006.
- 11 30. Demkowicz L, Rachowicz W, Devloo Ph. A fully automatic *hp*-adaptivity. *Journal of Scientific Computing* 2002;
17(1–3):127–155.
- 13 31. <http://www.mosetm.net/>.
- 15 32. Cong W, Shen H, Cong A, Wang Y, Wang G. Modeling photon propagation in biological tissues using a
generalized Delta–Eddington phase function. *Physical Review* 2007.
- 17 33. Jiang M, Zhou T, Cheng J, Cong W, Wang G. Image reconstruction method for bioluminescence tomography
from partial measurement. *Optics Express* 2007; **13**:11095–11116.

Q2

Q3

Author Queries Form

John Wiley

JOURNAL TITLE: CNM

11/11/2008

ARTICLE NO: 1202

Queries and / or remarks

Query No.	Details required	Author's response
Q1	Please provide DOI number if volume, page range are not available for Refs.[24,32].	
Q2	Please provide the place of proceedings for Ref.[29].	
Q3	Please provide further details.	

COPYRIGHT TRANSFER AGREEMENT



Date: _____ Contributor name: _____

Contributor address: _____

Manuscript number (Editorial office only): _____

Re: Manuscript entitled _____

_____ (the "Contribution")

for publication in _____ (the "Journal")

published by _____ ("Wiley-Blackwell").

Dear Contributor(s):

Thank you for submitting your Contribution for publication. In order to expedite the editing and publishing process and enable Wiley-Blackwell to disseminate your Contribution to the fullest extent, we need to have this Copyright Transfer Agreement signed and returned as directed in the Journal's instructions for authors as soon as possible. If the Contribution is not accepted for publication, or if the Contribution is subsequently rejected, this Agreement shall be null and void. **Publication cannot proceed without a signed copy of this Agreement.**

A. COPYRIGHT

1. The Contributor assigns to Wiley-Blackwell, during the full term of copyright and any extensions or renewals, all copyright in and to the Contribution, and all rights therein, including but not limited to the right to publish, republish, transmit, sell, distribute and otherwise use the Contribution in whole or in part in electronic and print editions of the Journal and in derivative works throughout the world, in all languages and in all media of expression now known or later developed, and to license or permit others to do so.

2. Reproduction, posting, transmission or other distribution or use of the final Contribution in whole or in part in any medium by the Contributor as permitted by this Agreement requires a citation to the Journal and an appropriate credit to Wiley-Blackwell as Publisher, and/or the Society if applicable, suitable in form and content as follows: (Title of Article, Author, Journal Title and Volume/Issue, Copyright © [year], copyright owner as specified in the Journal). Links to the final article on Wiley-Blackwell's website are encouraged where appropriate.

B. RETAINED RIGHTS

Notwithstanding the above, the Contributor or, if applicable, the Contributor's Employer, retains all proprietary rights other than copyright, such as patent rights, in any process, procedure or article of manufacture described in the Contribution.

C. PERMITTED USES BY CONTRIBUTOR

1. **Submitted Version.** Wiley-Blackwell licenses back the following rights to the Contributor in the version of the Contribution as originally submitted for publication:

a. After publication of the final article, the right to self-archive on the Contributor's personal website or in the Contributor's institution's/employer's institutional repository or archive. This right extends to both intranets and the Internet. The Contributor may not update the submission version or replace it with the published Contribution. The version posted must contain a legend as follows: This is the pre-peer reviewed version of the following article: FULL CITE, which has been published in final form at [Link to final article].

b. The right to transmit, print and share copies with colleagues.

2. **Accepted Version.** Re-use of the accepted and peer-reviewed (but not final) version of the Contribution shall be by separate agreement with Wiley-Blackwell. Wiley-Blackwell has agreements with certain funding agencies governing reuse of this version. The details of those relationships, and other offerings allowing open web use, are set forth at the following website: <http://www.wiley.com/go/funderstatement>. NIH grantees should check the box at the bottom of this document.

3. **Final Published Version.** Wiley-Blackwell hereby licenses back to the Contributor the following rights with respect to the final published version of the Contribution:

a. Copies for colleagues. The personal right of the Contributor only to send or transmit individual copies of the final published version in any format to colleagues upon their specific request provided no fee is charged, and further-provided that there is no systematic distribution of the Contribution, e.g. posting on a listserve, website or automated delivery.

b. Re-use in other publications. The right to re-use the final Contribution or parts thereof for any publication authored or edited by the Contributor (excluding journal articles) where such re-used material constitutes less than half of the total material in such publication. In such case, any modifications should be accurately noted.

c. Teaching duties. The right to include the Contribution in teaching or training duties at the Contributor's institution/place of employment including in course packs, e-reserves, presentation at professional conferences, in-house training, or distance learning. The Contribution may not be used in seminars outside of normal teaching obligations (e.g. commercial seminars). Electronic posting of the final published version in connection with teaching/training at the Contributor's institution/place of employment is permitted subject to the implementation of reasonable access control mechanisms, such as user name and password. Posting the final published version on the open Internet is not permitted.

d. Oral presentations. The right to make oral presentations based on the Contribution.

4. **Article Abstracts, Figures, Tables, Data Sets, Artwork and Selected Text (up to 250 words).**

a. Contributors may re-use unmodified abstracts for any non-commercial purpose. For on-line uses of the abstracts, Wiley-Blackwell encourages but does not require linking back to the final published versions.

b. Contributors may re-use figures, tables, data sets, artwork, and selected text up to 250 words from their Contributions, provided the following conditions are met:

(i) Full and accurate credit must be given to the Contribution.

(ii) Modifications to the figures, tables and data must be noted. Otherwise, no changes may be made.

(iii) The reuse may not be made for direct commercial purposes, or for financial consideration to the Contributor.

(iv) Nothing herein shall permit dual publication in violation of journal ethical practices.

D. CONTRIBUTIONS OWNED BY EMPLOYER

1. If the Contribution was written by the Contributor in the course of the Contributor's employment (as a "work-made-for-hire" in the course of employment), the Contribution is owned by the company/employer which must sign this Agreement (in addition to the Contributor's signature) in the space provided below. In such case, the company/employer hereby assigns to Wiley-Blackwell, during the full term of copyright, all copyright in and to the Contribution for the full term of copyright throughout the world as specified in paragraph A above.

2. In addition to the rights specified as retained in paragraph B above and the rights granted back to the Contributor pursuant to paragraph C above, Wiley-Blackwell hereby grants back, without charge, to such company/employer, its subsidiaries and divisions, the right to make copies of and distribute the final published Contribution internally in print format or electronically on the Company's internal network. Copies so used may not be resold or distributed externally. However the company/employer may include information and text from the Contribution as part of an information package included with software or other products offered for sale or license or included in patent applications. Posting of the final published Contribution by the institution on a public access website may only be done with Wiley-Blackwell's written permission, and payment of any applicable fee(s). Also, upon payment of Wiley-Blackwell's reprint fee, the institution may distribute print copies of the published Contribution externally.

E. GOVERNMENT CONTRACTS

In the case of a Contribution prepared under U.S. Government contract or grant, the U.S. Government may reproduce, without charge, all or portions of the Contribution and may authorize others to do so, for official U.S. Govern-

ment purposes only, if the U.S. Government contract or grant so requires. (U.S. Government, U.K. Government, and other government employees: see notes at end)

F. COPYRIGHT NOTICE

The Contributor and the company/employer agree that any and all copies of the final published version of the Contribution or any part thereof distributed or posted by them in print or electronic format as permitted herein will include the notice of copyright as stipulated in the Journal and a full citation to the Journal as published by Wiley-Blackwell.

G. CONTRIBUTOR'S REPRESENTATIONS

The Contributor represents that the Contribution is the Contributor's original work, all individuals identified as Contributors actually contributed to the Contribution, and all individuals who contributed are included. If the Contribution was prepared jointly, the Contributor agrees to inform the co-Contributors of the terms of this Agreement and to obtain their signature to this Agreement or their written permission to sign on their behalf. The Contribution is submitted only to this Journal and has not been published before. (If excerpts from copyrighted works owned by third parties are included, the Contributor will obtain written permission from the copyright owners for all uses as set forth in Wiley-Blackwell's permissions form or in the Journal's Instructions for Contributors, and show credit to the sources in the Contribution.) The Contributor also warrants that the Contribution contains no libelous or unlawful statements, does not infringe upon the rights (including without limitation the copyright, patent or trademark rights) or the privacy of others, or contain material or instructions that might cause harm or injury.

CHECK ONE BOX:

<input type="checkbox"/> Contributor-owned work ATTACH ADDITIONAL SIGNATURE PAGES AS NECESSARY	Contributor's signature _____	Date _____
	Type or print name and title _____	
	Co-contributor's signature _____	Date _____
	Type or print name and title _____	
<input type="checkbox"/> Company/Institution-owned work (made-for-hire in the course of employment)	Company or Institution (Employer-for-Hire) _____	Date _____
	Authorized signature of Employer _____	Date _____
<input type="checkbox"/> U.S. Government work	Note to U.S. Government Employees A contribution prepared by a U.S. federal government employee as part of the employee's official duties, or which is an official U.S. Government publication, is called a "U.S. Government work," and is in the public domain in the United States. In such case, the employee may cross out Paragraph A.1 but must sign (in the Contributor's signature line) and return this Agreement. If the Contribution was not prepared as part of the employee's duties or is not an official U.S. Government publication, it is not a U.S. Government work.	
<input type="checkbox"/> U.K. Government work (Crown Copyright)	Note to U.K. Government Employees The rights in a Contribution prepared by an employee of a U.K. government department, agency or other Crown body as part of his/her official duties, or which is an official government publication, belong to the Crown. U.K. government authors should submit a signed declaration form together with this Agreement. The form can be obtained via http://www.opsi.gov.uk/advice/crown-copyright/copyright-guidance/publication-of-articles-written-by-ministers-and-civil-servants.htm	
<input type="checkbox"/> Other Government work	Note to Non-U.S., Non-U.K. Government Employees If your status as a government employee legally prevents you from signing this Agreement, please contact the editorial office.	
<input type="checkbox"/> NIH Grantees	Note to NIH Grantees Pursuant to NIH mandate, Wiley-Blackwell will post the accepted version of Contributions authored by NIH grant-holders to PubMed Central upon acceptance. This accepted version will be made publicly available 12 months after publication. For further information, see www.wiley.com/go/nihmandate .	

WILEY AUTHOR DISCOUNT CARD

As a highly valued contributor to Wiley's publications, we would like to show our appreciation to you by offering a **unique 25% discount** off the published price of any of our books*.

To take advantage of this offer, all you need to do is apply for the **Wiley Author Discount Card** by completing the attached form and returning it to us at the following address:

The Database Group
John Wiley & Sons Ltd
The Atrium
Southern Gate
Chichester
West Sussex PO19 8SQ
UK

In the meantime, whenever you order books direct from us, simply quote promotional code **S001W** to take advantage of the 25% discount.

The newest and quickest way to order your books from us is via our new European website at:

<http://www.wileyeurope.com>

Key benefits to using the site and ordering online include:

- Real-time SECURE on-line ordering
- The most up-to-date search functionality to make browsing the catalogue easier
- Dedicated Author resource centre
- E-mail a friend
- Easy to use navigation
- Regular special offers
- Sign up for subject orientated e-mail alerts

So take advantage of this great offer, return your completed form today to receive your discount card.

Yours sincerely,



Verity Leaver
E-marketing and Database Manager

*TERMS AND CONDITIONS

This offer is exclusive to Wiley Authors, Editors, Contributors and Editorial Board Members in acquiring books (excluding encyclopaedias and major reference works) for their personal use. There must be no resale through any channel. The offer is subject to stock availability and cannot be applied retrospectively. This entitlement cannot be used in conjunction with any other special offer. Wiley reserves the right to amend the terms of the offer at any time.

REGISTRATION FORM FOR 25% BOOK DISCOUNT CARD

To enjoy your special discount, tell us your areas of interest and you will receive relevant catalogues or leaflets from which to select your books. Please indicate your specific subject areas below.

<p>Accounting []</p> <ul style="list-style-type: none"> • Public [] • Corporate [] 	<p>Architecture []</p>
<p>Chemistry []</p> <ul style="list-style-type: none"> • Analytical [] • Industrial/Safety [] • Organic [] • Inorganic [] • Polymer [] • Spectroscopy [] 	<p>Business/Management []</p>
<p>Encyclopedia/Reference []</p> <ul style="list-style-type: none"> • Business/Finance [] • Life Sciences [] • Medical Sciences [] • Physical Sciences [] • Technology [] 	<p>Computer Science []</p> <ul style="list-style-type: none"> • Database/Data Warehouse [] • Internet Business [] • Networking [] • Programming/Software Development [] • Object Technology []
<p>Earth & Environmental Science []</p>	<p>Engineering []</p> <ul style="list-style-type: none"> • Civil [] • Communications Technology [] • Electronic [] • Environmental [] • Industrial [] • Mechanical []
<p>Hospitality []</p>	<p>Finance/Investing []</p> <ul style="list-style-type: none"> • Economics [] • Institutional [] • Personal Finance []
<p>Genetics []</p> <ul style="list-style-type: none"> • Bioinformatics/Computational Biology [] • Proteomics [] • Genomics [] • Gene Mapping [] • Clinical Genetics [] 	<p>Life Science []</p>
<p>Medical Science []</p> <ul style="list-style-type: none"> • Cardiovascular [] • Diabetes [] • Endocrinology [] • Imaging [] • Obstetrics/Gynaecology [] • Oncology [] • Pharmacology [] • Psychiatry [] 	<p>Landscape Architecture []</p>
<p>Non-Profit []</p>	<p>Mathematics/Statistics []</p>
	<p>Manufacturing []</p>
	<p>Material Science []</p>
	<p>Psychology []</p> <ul style="list-style-type: none"> • Clinical [] • Forensic [] • Social & Personality [] • Health & Sport [] • Cognitive [] • Organizational [] • Developmental and Special Ed [] • Child Welfare [] • Self-Help []
	<p>Physics/Physical Science []</p>

I confirm that I am a Wiley Author/Editor/Contributor/Editorial Board Member of the following publications:

SIGNATURE:

PLEASE COMPLETE THE FOLLOWING DETAILS IN BLOCK CAPITALS:

TITLE AND NAME: (e.g. Mr, Mrs, Dr)

JOB TITLE:

DEPARTMENT:

COMPANY/INSTITUTION:

ADDRESS:

.....

.....

.....

TOWN/CITY:

COUNTY/STATE:

COUNTRY:

POSTCODE/ZIP CODE:

DAYTIME TEL:

FAX:

E-MAIL:

YOUR PERSONAL DATA

We, John Wiley & Sons Ltd, will use the information you have provided to fulfil your request. In addition, we would like to:

1. Use your information to keep you informed by post, e-mail or telephone of titles and offers of interest to you and available from us or other Wiley Group companies worldwide, and may supply your details to members of the Wiley Group for this purpose.
 Please tick the box if you do not wish to receive this information
2. Share your information with other carefully selected companies so that they may contact you by post, fax or e-mail with details of titles and offers that may be of interest to you.
 Please tick the box if you do not wish to receive this information.

If, at any time, you wish to stop receiving information, please contact the Database Group (databasegroup@wiley.co.uk) at John Wiley & Sons Ltd, The Atrium, Southern Gate, Chichester, West Sussex PO19 8SQ, UK.

E-MAIL ALERTING SERVICE

We offer an information service on our product ranges via e-mail. If you do not wish to receive information and offers from John Wiley companies worldwide via e-mail, please tick the box .

This offer is exclusive to Wiley Authors, Editors, Contributors and Editorial Board Members in acquiring books (excluding encyclopaedias and major reference works) for their personal use. There should be no resale through any channel. The offer is subject to stock availability and may not be applied retrospectively. This entitlement cannot be used in conjunction with any other special offer. Wiley reserves the right to vary the terms of the offer at any time.

Ref: S001W